

VIRSIN 423L

Inactivated vaccine against
Newcastle Disease
Egg Drop Syndrome
& Infectious Bronchitis

Notes:

1. Do not expose vaccine to direct sunlight.
2. In case of accidental human injection, refer to a physician immediately.

BATCH

ml

EXP. DATE



VIRSIN 423L

Inactivated vaccine against
Newcastle Disease
Egg Drop Syndrome
& Infectious Bronchitis

ND-LaSota strain
EDS 76-127 strain
IB-M41 strains

Oil emulsion

Manufactured by: **BIOVAC Ltd.**
30 Ha'llan st., Or-Akiva 3060000, ISRAEL

VIRSIN 423L

Inactivated vaccine against
Newcastle Disease
Egg Drop Syndrome
& Infectious Bronchitis

Administration Instructions:

1. Use by instructions of a veterinarian.
2. Vaccinate only healthy flocks, apply stringent aseptic procedures.
3. Warm vaccine to room temperature (20°-25°C), just before use.
4. Shake **well** before and during use.
5. For future layers use at least 2-4 weeks prior to onset of egg production.
6. **Dosage:** 0.5 ml per bird, S.C.
7. Discard remained unused vaccine.

Storage:

Store at 2°-8°C.

DO NOT FREEZE

For veterinary use only

0500321-02