



VIR 101

Infectious Laryngotracheitis (ILT)

Freeze Dried Live Vaccine – Samberg strain

COMPOSITION

Each dose contains:

Attenuated virus of Infectious Laryngotracheitis Samberg strain, at least $10^{3.0}$ EID₅₀.

INDICATIONS

Immunisation of chickens against Infectious Laryngotracheitis.

ADMINISTRATION AND DOSAGE

Use by instructions of a veterinarian. Reconstitute immediately prior to use.

Cloacal brush

- Aseptically transfer 5 ml of BIOVAC'S ILT DILUENT into the vaccine vial.
- Shake well for complete reconstitution.
- Return the dissolved vaccine into the diluent vial containing the remaining diluent.
- Dip the enclosed brush in the diluted vaccine and brush the upper cloacal mucosa 3-4 times.
- **Takes** - The flock (20-25 birds) should be examined 4-5 days after vaccination for evidence of takes. Positive reaction is a slight swelling and red cherry coloration of the cloacal mucosa. At least 85% of positive birds is evidence of successful vaccination.

Drinking water

- Make sure that drinking water is free of any traces of disinfectants.
- Deprive the birds of drinking water two hours before vaccination.
- Prepare adequate volume of drinking water to be consumed within one to two hours.
- Protect the dissolved vaccine from chlorine by adding BIOVAC'S Blue Neutrachlor® (following the instructions).
- Stir well after mixing and allow the solution to stand for 20 min before adding the vaccine.
- Reconstitute the vaccine with antiseptic or disinfectant-free drinking water.

- Mix the reconstituted vaccine well in the final disinfectant-free drinking water volume.
- Let the birds consume all the vaccine-water volume before renewing the water supply.

CONTRA-INDICATIONS, WARNINGS

Future layers and breeders should not be vaccinated before 3 weeks of age by the cloacal brush method. Do not vaccinate bird in lay.

PRECAUTIONS

- Vaccinate healthy chickens only.
- Use the entire contents of the vial when first opened.
- Destroy unused contents and decontaminate empty vials before discarding them.
- Keep vaccination records including vaccine batch number.

For veterinary use only.

Keep out of reach and sight of children.

WITHDRAWAL PERIOD

Zero days.

STORAGE

Vaccine: Store at 2-8°C, protected from direct sunlight.

Diluent: Store in a cool and dark place. Store below 25°C.

PRESENTATION

Vaccine: Each box contains: 10 vials of 500 or 1,000 doses.

Diluent: Each box contains 10 bottles of 20 ml for 500 doses, or 40 ml for 1,000 doses.

Manufacturer and registration holder:

BIOVAC Ltd.

30 Ha' Ilan St., Or- Akiva
3060000, Israel.

T. 972-4-6266772, F. 972-4-6260912

E-mail: vireo@biovac.co.il

0500219-02