



VIR- 102

Fowl Pox

Freeze Dried Live Vaccine

Composition

Each dose contains:

Attenuated virus of Fowl Pox,

At least 10^{3.0} EID₅₀.

Indications

Immunization of chickens & turkeys against Fowl Pox via wing-web prick method.

Administration and dosage

Use by instruction of a veterinarian.

Reconstitution: Do not reconstitute the vaccine until ready to begin vaccination. Reconstitute the vaccine with Biovac's sterile diluent for VIR 102 (Fowl Pox). For manual prick application, use 5 ml of diluent, per 1000 doses.

For automatic Fowl Pox syringe, use 2.5 ml diluent, per 1000 doses. The vaccine is applied by the wing web method. Dip the enclosed lancet into the diluted vaccine and vaccinate each bird through the skin in the wing- web area.

The lancet should be dipped in the vaccine before each prick.

Turkeys – a double dose should be applied.

This vaccine may be used with an automatic fowl pox syringe. Ensure that syringe was properly sterilized.

Check for Takes – Examine and record takes on 10% of the flock 6-7 days following vaccination. A scab or swelling of the skin at the point of vaccination is indicative for a satisfactory vaccination.

At least 90% positive birds is an evidence of successful vaccination. If the percentage of positive birds is lower, revaccinate the flock.

Precautions

- Vaccinate healthy birds only.
- For reconstitution, as referred above, use only Biovac's sterile diluent for VIR 102 (Fowl Pox).
- Use entire contents of the vial when first opened.
- Destroy unused contents and decontaminate empty vials before discarding them.
- Keep vaccination records including vaccine batch number.

Withdrawal period

None.

Storage

Vaccine: Store at 2°- 8°C, protected from direct sunlight.

Diluent: Store at a cool and dark place.

Presentation

Vaccine: Each box contains: 10 vials of 1000 doses each.

Diluent: Each box contains: 10 vials of 5 ml each.

BIOVAC Ltd.

30 Ha' Ilan St., Or- Akiva
3060000, Israel.

T. 972-4-6266772, F. 972-4-6260912

E-mail: vireo@biovac.co.il