

## VIRSIN 336

Inactivated vaccine against  
**Infectious Coryza**

### Notes:

1. Do not expose vaccine to direct sunlight.
2. In case of accidental human injection, refer to a physician immediately.

BATCH

ml

EXP. DATE



## VIRSIN 336

Inactivated vaccine against  
**Infectious Coryza**  
**Avibacterium paragallinarum**  
**serotypes A,B&C**

Oil emulsion

Manufactured by: **BIOVAC Ltd.**  
30 Ha'llan St., Or-Akiva 3060000, ISRAEL

## VIRSIN 336

Inactivated vaccine against  
**Infectious Coryza**

### Administration Instructions:

1. Use by instructions of a veterinarian.
2. Vaccinate only healthy flocks, apply stringent aseptic procedure.
3. Warm vaccine to room temperature (20°-25°C), just before use.
4. Shake **well** before and during use.
5. First vaccination should be done at 6-8 weeks of age. For sufficient protection vaccinate birds twice at 3-4 weeks intervals.
6. **Dosage:** 0.5 ml per bird, S.C.
7. Discard remained unused vaccine.

### Storage:

Store at 2°-8°C.

### DO NOT FREEZE

**For animal use only**

**Keep out of sight and reach of children**

0500334-01