

VIRSIN 539L

Inactivated vaccine against
Newcastle Disease
Infectious Bronchitis
Infectious Bursal Disease
& Viral Arthritis (REO)

Notes:

1. Do not expose vaccine to direct sunlight.
2. In case of injecting oil emulsion to human, see a physician immediately.

956105/05

BATCH

ml

EXP. DATE



VIRSIN 539L

Inactivated vaccine against
Newcastle Disease
Infectious Bronchitis
Infectious Bursal Disease (Gumboro)
& Viral Arthritis (REO)

ND-LaSota strain
IB-M41 strain
IBD-FS strain
Reo-S1133+641 strains
Oil emulsion

Manufactured by: **BIOVAC** Ltd.
30 Ha'ilan st. Or Akiva 3060000, ISRAEL

VIRSIN 539L

Inactivated vaccine against
Newcastle Disease
Infectious Bronchitis
Infectious Bursal Disease
& Viral Arthritis (REO)

Administration Instructions:

1. Use by instructions of a veterinarian.
2. Vaccinate only a healthy flock, apply stringent aseptic procedure.
3. Warm vaccine to room temperature (20°-25°C), just before use.
4. Shake **well** before and during use.
5. For future layers use at least 2-4 weeks prior to the onset of egg production.
6. **Dosage:** 0.5 ml per bird. S.C.
7. Discard remained unused vaccine.

For veterinary use only

Storage:

Store at 2°-8°C.

DO NOT FREEZE